

Uniqueness of a convex sum of products of projectors

K. A. Kirkpatrick*

New Mexico Highlands University, Las Vegas, New Mexico 87701

Relative to a given factoring of the Hilbert space, the decomposition of an operator into a convex sum of correlated products of pairs of distinct 1-projectors, one set of projectors linearly independent, is unique.

Utilizing the Tridecompositional Uniqueness Theorem of Elby and Bub¹, I establish the uniqueness, relative to a given factoring of the Hilbert space, of a decomposition of a state operator into a convex sum of correlated products of pairs of distinct 1-projectors,

$$\rho = \sum_j w_j |a_j\rangle\langle a_j| \otimes |b_j\rangle\langle b_j|, \quad (1)$$

one set of projectors linearly independent.

In the appendix, I present a slightly strengthened version and simplified proof of the Tridecompositional Uniqueness Theorem.

For the remainder of this paper I use the notation $|a_j b_k\rangle$ for the direct product $|a_j\rangle \otimes |b_k\rangle$.

I. PRELIMINARIES

All vectors are normalized.

Definition 1 $|a\rangle$ and $|b\rangle$ are collinear iff $|a\rangle = e^{i\alpha} |b\rangle$, $\alpha \in \mathcal{R}$; we denote this $|a\rangle \parallel |b\rangle$.

Definition 2 The set $\{|a_j\rangle\}$ is non-collinear iff no pair of the set is collinear.

Definition 3 ρ is an operator on $\mathcal{H}_1 \otimes \mathcal{H}_2$. The null space of ρ on $\mathcal{H}_\alpha$ is $\mathcal{N}_\alpha \doteq \{|\phi\rangle \in \mathcal{H}_\alpha \mid \rho|\phi\rangle = 0\}$ ($\alpha \in \{1, 2\}$).

Lemma 1 With sets $\{|a_j\rangle \in \mathcal{H}_1\}$, $\{|b_j\rangle \in \mathcal{H}_2\}$, and $\{w_j > 0\}$, $j \in \{1..N\}$, and the operator $\rho = \sum_j w_j |a_j b_j\rangle\langle a_j b_j|$, the set $\{|a_j\rangle\}$ spans $\mathcal{G}_1 \doteq (\mathcal{N}_1)^\perp$ and the set $\{|b_j\rangle\}$ spans $\mathcal{G}_2 \doteq (\mathcal{N}_2)^\perp$.

Proof: For $|\phi\rangle \in \mathcal{N}_1$ and any $|\beta\rangle \in \mathcal{H}_2$, $\langle\phi\beta|\rho|\phi\beta\rangle = 0 = \sum_j w_j |\langle a_j|\phi\rangle|^2 |\langle b_j|\beta\rangle|^2$; thus $\langle a_j|\phi\rangle = 0 \forall |\phi\rangle \in \mathcal{N}_1$, so $|a_j\rangle \in (\mathcal{N}_1)^\perp = \mathcal{G}_1$. If $\{|a_j\rangle\}$ does not span $\mathcal{G}_1$, there is a vector in $\mathcal{G}_1$ orthogonal to $\{|a_j\rangle\}$; but any such vector is annihilated by ρ and is thus in $\mathcal{N}_1$, a contradiction. $\square$

The following result appears in Ref. 2, in the midst of the proof of another theorem:

Lemma 2 $|\Psi\rangle$ and $|\Phi\rangle$ are vectors in $\mathcal{H}_1 \otimes \mathcal{H}_2$. If $\text{Tr}_2\{|\Psi\rangle\langle\Psi|\} = \text{Tr}_2\{|\Phi\rangle\langle\Phi|\}$, then there exists a unitary transformation $\mathbf{U}$ on $\mathcal{H}_2$ such that $|\Psi\rangle = (\mathbf{I} \otimes \mathbf{U})|\Phi\rangle$.

II. THE UNIQUENESS THEOREM

Theorem 1 *With non-collinear sets $\{|a_j\rangle \in \mathcal{H}_1\}$ and $\{|b_j\rangle \in \mathcal{H}_2\}$, $j \in \{1..n\}$, one set linearly independent, and with non-collinear sets $\{|A_k\rangle \in \mathcal{H}_1\}$ and $\{|B_k\rangle \in \mathcal{H}_2\}$, $k \in \{1..N\}$, one set linearly independent, and with sets $\{w_j > 0\}$ and $\{W_k > 0\}$, if*

$$\sum_{j=1}^n w_j |a_j b_j\rangle \langle a_j b_j| = \sum_{k=1}^N W_k |A_k B_k\rangle \langle A_k B_k|,$$

then $N = n$, and, for all $j \in \{1..n\}$,

$$|A_j\rangle \parallel |a_{\pi(j)}\rangle, \quad |B_j\rangle \parallel |b_{\pi(j)}\rangle, \quad \text{and } W_j = w_{\pi(j)},$$

with $\pi(\cdot)$ a permutation function on $\{1..n\}$.

Proof: Call the operator ρ . Apply Lemma 1, with $d_1 \doteq \dim \mathcal{G}_1$ and $d_2 \doteq \dim \mathcal{G}_2$, and recall: A set of m vectors spans a space of dimension $d \leq m$; $d = m$ iff the vectors are linearly independent. Without loss of generality we take the set $\{|a_j\rangle\}$, which spans $\mathcal{G}_1$, to be linearly independent; thus $n = d_1$. Either $\{|A_k\rangle\}$ or $\{|B_k\rangle\}$ must be linearly independent; in either case, $N = n$: If $\{|A_k\rangle\}$ is linearly independent, then $N = d_1 = n$. On the other hand, if $\{|B_k\rangle\}$ is linearly independent, then $N = d_2$; the n vectors $\{|b_j\rangle\}$ must span $\mathcal{G}_2$, hence $n \geq d_2 = N$. Similarly, the N vectors $\{|A_k\rangle\}$ must span $\mathcal{G}_1$, hence $N \geq d_1 = n$, thus $N = n$.

Introduce a third Hilbert space $\mathcal{H}_3$, with $\dim \mathcal{H}_3 \geq n$; $\{|c_j\rangle\}$ and $\{|C_j\rangle\}$ are orthonormal bases of $\mathcal{H}_3$. Construct the two vectors

$$|\psi\rangle = \sum_{j=1}^n \sqrt{w_j} |a_j b_j c_j\rangle \quad \text{and} \quad |\Psi\rangle = \sum_{j=1}^n \sqrt{W_j} |A_j B_j C_j\rangle;$$

clearly, $\rho = \text{Tr}_3\{|\psi\rangle\langle\psi|\} = \text{Tr}_3\{|\Psi\rangle\langle\Psi|\}$. By Lemma 2, there exists a unitary transformation $\mathbf{U}$ on $\mathcal{H}_3$ such that $|\psi\rangle = (\mathbf{1} \otimes \mathbf{1} \otimes \mathbf{U})|\Psi\rangle$; defining $|D_j\rangle \doteq \mathbf{U}|C_j\rangle$, we have

$$\sum_{j=1}^n \sqrt{w_j} |a_j b_j c_j\rangle = \sum_{j=1}^n \sqrt{W_j} |A_j B_j D_j\rangle,$$

to which we apply Theorem A. □

III. DISCUSSION

“Uniqueness” is relative to the identification of system and apparatus

Elby and Bub claim that Eq. (1) “suffers from a version of the basis degeneracy problem.” For example, with $\langle a_1 | a_2 \rangle = \langle b_1 | b_2 \rangle = 0$, the sum-of-products expression

$$\rho = \frac{1}{2} |a_1 b_1\rangle \langle a_1 b_1| + \frac{1}{2} |a_2 b_2\rangle \langle a_2 b_2| \quad (2)$$

(which, according to Theorem 1, is unique) is the diagonalization of a degenerate Hermitian operator (with eigenvalues $1/2$ twice, and 0 twice). The eigenvectors may be

taken to be $|a_1 b_1\rangle$, $|a_2 b_2\rangle$, $|a_1 b_2\rangle$ and $|a_2 b_1\rangle$ — products of vectors taken pairwise from $\mathcal{H}_1$ and $\mathcal{H}_2$. Because of this degeneracy, we can rotate the eigenvectors into $|q_{1,2}\rangle = 2^{-1/2}(|a_1 b_1\rangle \pm |a_2 b_2\rangle)$, $|q_3\rangle = |a_1 b_2\rangle$, and $|q_4\rangle = |a_2 b_1\rangle$. Then

$$\rho = \frac{1}{2} |q_1\rangle\langle q_1| + \frac{1}{2} |q_2\rangle\langle q_2|; \quad (3)$$

“the pointer reading loses its ‘special’ status.”

This argument is flawed — after all, the same claim may be made against the tridecompositional uniqueness theorem itself:

$$|\Psi\rangle = \frac{1}{\sqrt{2}}(|a_1 b_1 c_1\rangle + |a_2 b_2 c_2\rangle) = \frac{1}{\sqrt{2}}(|q_1 d_1\rangle + |q_2 d_2\rangle), \quad (4)$$

with $|d_{1,2}\rangle = 2^{-1/2}(|c_1\rangle \pm |c_2\rangle)$. Eq. (4) is no more a counterexample to the tridecompositional uniqueness theorem than Eq. (3) is a counterexample to Theorem 1, and for the same reason: the “special” nature of a pointer basis is based on the uniqueness of the form of the decomposition in Eq. (1), which in turn is based on a particular identification of system and apparatus. One cannot speak of the “pointer basis” without having settled on the “pointer” — the apparatus — thus having already specified the factor spaces.

Only having chosen a fixed identification of the subsystems (and the associated factoring of the space) may either of these uniqueness theorems then be applied.

APPENDIX. THE TRIDECOMPOSITIONAL UNIQUENESS THEOREM

This version of the Tridecompositional Uniqueness Theorem¹ avoids two assumptions of the original: that the linearly dependent set is in the same space in each expansion, and that the expansions each have the same number of terms. The proof here is similar to that of Ref. 1, but is considerably shorter and, perhaps, clearer.

Definition 4 $|\Psi\rangle$ is factorable in $\mathcal{H}_1 \otimes \mathcal{H}_2$ iff there exist $|\alpha\rangle \in \mathcal{H}_1$ and $|\beta\rangle \in \mathcal{H}_2$ such that $|\Psi\rangle = |\alpha\beta\rangle$.

Lemma A (Similar to Lemma 1 of Ref. 1) *With the set $\{|a_j\rangle \in \mathcal{H}_1\}$ linearly independent and the set $\{|b_j\rangle \in \mathcal{H}_2\}$ non-collinear, $|\Psi\rangle = \sum_j s_j |a_j b_j\rangle$ is factorable in $\mathcal{H}_1 \otimes \mathcal{H}_2$ iff the set $\{s_j \in \mathcal{C}\}$ contains exactly one non-zero element.*

Proof: Let $|\Psi\rangle = |\alpha\beta\rangle$, with $|\alpha\rangle \in \mathcal{H}_1$ and $|\beta\rangle \in \mathcal{H}_2$. Expand $|\alpha\rangle = \sum_j a_j |a_j\rangle$, so $|\Psi\rangle = \sum_j a_j |a_j\beta\rangle$; the set $\{|a_j\rangle\}$ is linear independent, so, for each j , $a_j |\beta\rangle = s_j |b_j\rangle$. For every $s_j \neq 0$, $|b_j\rangle \parallel |\beta\rangle$. If more than one $s_j \neq 0$, $\{|b_j\rangle\}$ is not non-collinear, contrary to hypothesis, contradicting the assumption of factorability. The converse is obvious. $\square$

Theorem A (Tridecompositional uniqueness) *With non-collinear sets $\{|a_j\rangle \in \mathcal{H}_1\}$, $\{|b_j\rangle \in \mathcal{H}_2\}$, and $\{|c_j\rangle \in \mathcal{H}_3\}$, $j \in \{1..n\}$, two sets linearly independent, and non-collinear sets $\{|A_k\rangle \in \mathcal{H}_1\}$, $\{|B_k\rangle \in \mathcal{H}_2\}$, and $\{|C_k\rangle \in \mathcal{H}_3\}$, $k \in \{1..N\}$, two sets linearly independent, and sets $\{\phi_j \in \mathcal{C} \mid \phi_j \neq 0\}$ and $\{\varphi_k \in \mathcal{C} \mid \varphi_k \neq 0\}$, if*

$$\sum_{j=1}^n \phi_j |a_j b_j c_j\rangle = \sum_{k=1}^N \varphi_k |A_k B_k C_k\rangle,$$

then $N = n$, and, for all $j \in \{1..n\}$,

$$|A_j\rangle \parallel |a_{\pi(j)}\rangle, \quad |B_j\rangle \parallel |b_{\pi(j)}\rangle, \quad |C_j\rangle \parallel |c_{\pi(j)}\rangle, \quad \text{and} \quad |\varphi_j| = |\phi_{\pi(j)}|,$$

with $\pi(\cdot)$ a permutation function on $\{1..n\}$.

Proof: Take $\{|C_k\rangle\}$ and $\{|c_j\rangle\}$ to be linearly independent (with no loss of generality: in each expansion, two of the three sets are linearly independent, requiring coincidence in at least one space). These sets must span the same subspace of $\mathcal{H}_3$; thus $N = n$. Expand $|c_j\rangle = \sum_k \gamma_{jk} |C_k\rangle$; then $\varphi_k |A_k B_k\rangle = \sum_j \phi_j \gamma_{jk} |a_j b_j\rangle$. For each k , Lemma A requires $\gamma_{jk} = 0$ for all but one j ; define the function $\pi: \{1..n\} \rightarrow \{1..n\}$ by the relation $\gamma_{\pi(k)k} \neq 0$. We have $|c_{\pi(k)}\rangle = \gamma_{\pi(k)k} |C_k\rangle$, so $|c_{\pi(k)}\rangle \parallel |C_k\rangle$ (and normalization requires $|\gamma_{\pi(k)k}| = 1$). Because the set $\{|C_k\rangle\}$ is non-collinear, $\pi(\cdot)$ must be 1 : 1, i.e., a permutation function on $\{1..n\}$. We also have $\varphi_k |A_k B_k\rangle = \phi_{\pi(k)} \gamma_{\pi(k)k} |a_{\pi(k)} b_{\pi(k)}\rangle$, so $|a_{\pi(k)}\rangle \parallel |A_k\rangle$ and $|b_{\pi(k)}\rangle \parallel |B_k\rangle$; normalization requires $|\varphi_k| = |\phi_{\pi(k)}|$. $\square$

* E-mail: kirkpatrick@physics.nmhu.edu

¹ A. Elby and J. Bub, "Triorthogonal Uniqueness Theorem and its relevance to the interpretation of quantum mechanics," Phys. Rev. A **49**(5), 4213–4216 (1994).

² N. D. Mermin, "What do these correlations know about reality? Nonlocality and the absurd," Found. Phys. **29**(4), 571–587 (1999), quant-ph/9807055.